

ITERATIVE METHODS FOR TIKHONOV REGULARIZATION

D. Calvetti^a and L. Reichel^b

^aDepartment of Mathematics
Case Western Reserve University
Cleveland, OH 44122
dxc57@po.cwru.edu

^bDepartment of Mathematical Sciences
Kent State University
Kent, OH 44242
reichel@math.kent.edu

Tikhonov regularization is one of the most popular regularization techniques for the solution of linear ill-posed problems. Large-scale ill-posed problems have to be solved by iterative techniques, and we review available and present new iterative methods. We consider both the cases when the norm of the error in the available observations is known and when it is not.

References

- [1]D. Calvetti, S. Morigi, L. Reichel, and F. Sgallari , “Tikhonov regularization and the L-curve for large, discrete ill-posed problems,” *J. Comput. Appl. Math.* , v. 123, p. 423-446, 2000.
- [2]D. Calvetti, S. Morigi, L. Reichel, and F. Sgallari , “An L-ribbon for large underdetermined linear discrete ill-posed problems,” *Numer. Algorithms* , v. 25, p. 89-107, 2000.
- [3]D. Calvetti, P. C. Hansen , and L. Reichel, “L-curve curvature bounds via Lanczos bidiagonalization,” *Elec. Trans. Numer. Anal.* , v. 14, p. 20-35, 2002.
- [4]D. Calvetti, and L. Reichel, “Tikhonov regularization of large linear problems,” *BIT* , to appear.